



LAB #: H160415-2375-1
 PATIENT: Anna M. Salanti
 ID: SALANTI-A-00001
 SEX: Female
 AGE: 64

CLIENT #: 26339
 DOCTOR:
 Cellphysics.Org
 1537 Jarrettsville Rd
 Jarrettsville, MD 21084 U.S.A.

Toxic & Essential Elements; Hair

TOXIC METALS				PERCENTILE	
		RESULT µg/g	REFERENCE INTERVAL	68 th	95 th
Aluminum (Al)		3.7	< 12		
Antimony (Sb)		0.084	< 0.060		
Arsenic (As)		0.063	< 0.090		
Barium (Ba)		2.1	< 2.0		
Beryllium (Be)		< 0.01	< 0.020		
Bismuth (Bi)		0.007	< 2.0		
Cadmium (Cd)		0.036	< 0.050		
Lead (Pb)		0.32	< 1.0		
Mercury (Hg)		0.25	< 0.80		
Platinum (Pt)		< 0.003	< 0.005		
Thallium (Tl)		0.001	< 0.002		
Thorium (Th)		< 0.001	< 0.002		
Uranium (U)		0.005	< 0.060		
Nickel (Ni)		0.15	< 0.40		
Silver (Ag)		0.18	< 0.10		
Tin (Sn)		0.05	< 0.30		
Titanium (Ti)		1.3	< 1.3		
Total Toxic Representation					

ESSENTIAL AND OTHER ELEMENTS				PERCENTILE				
		RESULT µg/g	REFERENCE INTERVAL	2.5 th	16 th	50 th	84 th	97.5 th
Calcium (Ca)		1520	475- 1500					
Magnesium (Mg)		260	45- 180					
Sodium (Na)		200	80- 450					
Potassium (K)		40	28- 160					
Copper (Cu)		23	11- 30					
Zinc (Zn)		200	130- 200					
Manganese (Mn)		1.2	0.15- 0.65					
Chromium (Cr)		0.32	0.40- 0.65					
Vanadium (V)		0.013	0.018- 0.065					
Molybdenum (Mo)		0.052	0.040- 0.10					
Boron (B)		3.5	0.40- 4.0					
Iodine (I)		2.2	0.25- 1.8					
Lithium (Li)		0.12	0.008- 0.030					
Phosphorus (P)		687	250- 500					
Selenium (Se)		1.1	0.80- 1.3					
Strontium (Sr)		11	1.0- 8.0					
Sulfur (S)		44700	42000- 48000					
Cobalt (Co)		0.014	0.006- 0.035					
Iron (Fe)		7.8	7.0- 16					
Germanium (Ge)		0.036	0.030- 0.040					
Rubidium (Rb)		0.042	0.030- 0.25					
Zirconium (Zr)		< 0.007	0.040- 1.0					

SPECIMEN DATA		RATIOS		
COMMENTS:		ELEMENTS	RATIOS	RANGE
Date Collected: 04/05/2016		Ca/Mg	5.85	4- 30
Date Received: 04/15/2016		Ca/P	2.21	1- 12
Date Completed: 04/20/2016		Na/K	5	0.5- 10
Methodology: ICP/MS		Zn/Cu	8.7	4- 20
Sample Size: 0.201 g		Zn/Cd	> 999	> 800
Sample Type: Pubic				
Hair Color: Gray				
Treatment:				
Shampoo: Sappo Hill				

HAIR ELEMENTS REPORT INTRODUCTION

Hair is an excretory tissue for essential, nonessential and potentially toxic elements. In general, the amount of an element that is irreversibly incorporated into growing hair is proportional to the level of the element in other body tissues. Therefore, hair elements analysis provides an indirect screening test for physiological excess, deficiency or maldistribution of elements in the body. Clinical research indicates that hair levels of specific elements, particularly potentially toxic elements such as cadmium, mercury, lead and arsenic, are highly correlated with pathological disorders. For such elements, levels in hair may be more indicative of body stores than the levels in blood and urine.

All screening tests have limitations that must be taken into consideration. The correlation between hair element levels and physiological disorders is determined by numerous factors. Individual variability and compensatory mechanisms are major factors that affect the relationship between the distribution of elements in hair and symptoms and pathological conditions. It is also very important to keep in mind that scalp hair is vulnerable to external contamination of elements by exposure to hair treatments and products. Likewise, some hair treatments (e.g. permanent solutions, dyes, and bleach) can strip hair of endogenously acquired elements and result in false low values. Careful consideration of the limitations must be made in the interpretation of results of hair analysis. The data provided should be considered in conjunction with symptomology, diet analysis, occupation and lifestyle, physical examination and the results of other analytical laboratory tests.

Caution: The contents of this report are not intended to be diagnostic and the physician using this information is cautioned against treatment based solely on the results of this screening test. For example, copper supplementation based upon a result of low hair copper is contraindicated in patients afflicted with Wilson's Disease.

Pubic Hair Specimens

Pubic hair and scalp hair are very different tissues with respect to protein and chemical composition, and rate of growth. The levels of most nutrients elements in pubic and scalp hair for a given individual are typically quite different. Although we do have reference ranges for nutrient elements in pubic hair specimens, there is a lack of clinical data to support sound interpretation at this time. For the potentially toxic elements, however, there appears to be good correlation between scalp and pubic hair. Some clinicians utilize pubic hair for toxic element analyze, (a) to confirm results from scalp hair, and/or (b) when scalp hair has been recently treated with dye or permanent and bleaching reagents.

Antimony High

Hair is a preferred tissue for analysis of Antimony (Sb) exposure and body burden. Elevated hair Sb levels have been noted as long as a year after exposure.

Sb is a nonessential element that is chemically similar to but less toxic than arsenic. Food and smoking are the usual sources of Sb. Thus cigarette smoke can externally contaminate hair, as well as contribute to uptake via inhalation. Gunpowder (ammunition) often contains Sb. Firearm enthusiasts often have elevated levels of Sb in hair. Other possible sources are textile industry,

metal alloys, and some antihelminthic and antiprotozoic drugs. Sb is also used in the manufacture of paints, glass, ceramics, solder, batteries, bearing metals, semiconductors and fire retardant fabrics.

Like arsenic, Sb has a high affinity for sulfhydryl groups on many enzymes. Sb is conjugated with glutathione and excreted in urine and feces. Therefore, excessive exposure to Sb has the potential to deplete intracellular glutathione pools.

Early signs of Sb excess include: fatigue, muscle weakness, myopathy, nausea, low back pain, headache, and metallic taste. Later symptoms include hemolytic anemia, myoglobinuria, hematuria and renal failure. Transdermal absorption can lead to "antimony spots" which resemble chicken pox. Respiratory tissue irritation may result from inhalation of Sb particles or dust.

A confirmatory test for recent or current exposure is the measurement of Sb in the urine or whole blood. Comparison of pre and post provocation (DMPS, DMSA, Ca-EDTA) urine Sb levels provides an estimate of net retention (body burden) of Sb.

Silver High

Hair Silver (Ag) levels have been found to reflect environmental exposure to the element. However, hair is commonly contaminated with Ag from hair treatments such as permanents, dyes, and bleaches.

Ag is not an essential element and is of relatively low toxicity. However, some Ag salts are very toxic.

Sources of Ag include seafood, metal and chemical processing industries, photographic processes, jewelry making (especially soldering), effluents from coal fired power plants and colloidal silver products.

The bacteriostatic properties of Ag have been long recognized and Ag has been used extensively for medicinal purposes; particularly in the treatment of burns. There is much controversy over the long term safety of consumption of colloidal silver. Very high intake of colloidal silver has been reported to give rise to tumors in the liver and spleen of animals (Metals in Clinical and Analytical Chemistry, eds. Seiler, Segel and Segel, 1994). However, these data may not have relevance to the effects of chronic, low level consumption by humans.

Magnesium High

Magnesium (Mg) is an essential element with both electrolyte and enzyme-activator functions. However, neither of these functions takes place in hair. Body excess of Mg is rare but may occur from excessive oral or parenteral supplementation or as a result of renal damage or insufficiency.

If one rules out external contamination of hair as a result of recent hair treatment, elevated hair Mg is more likely to indicate maldistribution of the element. Physiological Mg dysfunction may or may not be present. Maldistribution of Mg can occur as a result of chronic emotional or physical stress, toxic metal or chemical exposure, physiological imbalance of calcium and phosphorus, bone mineral depletion, and renal insufficiency with poor clearance of Mg (and other metabolites). Elevated hair Mg has been correlated with hypoglycemia and an inappropriately low ratio of dietary Ca : P.

Mg status can be difficult to assess; whole blood and packed blood red cell Mg levels are more indicative than serum/plasma levels. Amino acid analysis can be helpful in showing rate-limited steps that are Mg-dependent (e.g. phosphorylations).

Copper Normal

Hair Copper (Cu) levels are usually indicative of body status, except that exogenous contamination may occur giving a false normal (or false high). Common sources of contamination include: permanent solutions, dyes, bleaches, and swimming pools/hot tubs in which Cu compounds have been used as algacides.

Cu is an essential element that activates specific enzymes. Erythrocyte superoxide dismutase (SOD) is a Cu (and zinc) dependent enzyme; lysyl oxidase which catalyzes crosslinking of collagen is another Cu dependent enzyme. Adrenal catecholamine synthesis is Cu dependent, because the enzyme dopamine beta-hydroxylase, which catalyzes formation of norepinephrine from dopamine, requires Cu.

If hair Cu is in the normal range, this usually means tissue levels are in the normal range. However, under circumstances of contamination, a real Cu deficit could appear as a (false) normal. If symptoms of Cu deficiency are present, a whole blood or red blood cell elements analysis can be performed for confirmation of Cu status.

Manganese High

Hair Manganese (Mn) levels generally reflect exposure to Mn, but external contamination can influence hair Mn. High hair Mn can be an artifact of contamination from: permanent solutions, dyes, bleaches, and well water (containing high Mn). These possibilities should be considered and ruled out before proceeding with therapies to alleviate excess Mn.

Mn is an essential element which is involved in the activation of many important enzymes. However, Mn excess is postulated to result in glutathionyl radical formation, reduction of the free glutathione pool, and increased exposure of adrenal catecholamines (e.g. dopamine) to free radical damage. Excess Mn causes degeneration of melanin-pigmented dopaminergic neurons which results in abnormally low levels of serotonin and dopamine in the brain. This is hypothesized to be a reason behind the neurotoxic effects attributed to Mn overload.

The brain is particularly affected by Mn excess. Symptoms or conditions consistent with excessive Mn include: disorientation, memory loss, anxiety, hypotonia, abnormal gait, emotional instability, and bipolar-like behaviors (laughing and crying), aberrant or violent behaviors, and tremor or Parkinson-like symptoms.

Causes of Mn excess include: occupational or environmental exposures, contaminated teas, MMT (gasoline additive), coal-fired power plants, contaminated drinking water, some street drugs (cocaine products), and smoking. Conditions predisposing to Mn excess are: iron or calcium deficiency, chronic infection, and impaired liver function (e.g. biliary obstruction) or disease. Mn excess is occasionally associated with alcoholism.

Confirmatory tests for Mn excess include whole blood and a comparison of urine Mn pre- and post Ca-EDTA.

Chromium Low

Hair Chromium (Cr) is a good indicator of tissue levels and may provide a better indication of status than do urine or blood plasma/serum (Nielsen, F.H. In Modern Nutrition on Health and Disease; 8th Edition, 1994. Ed. Shils, Olson and Shike. Lea and Febiger, Philadelphia). Hair Cr is seldom affected by permanent solutions, dyes and bleaches.

Cr (trivalent) is generally accepted as an essential trace element that is required for maintenance of normal glucose and cholesterol levels; it potentiates insulin function, i.e., as a part of "glucose tolerance factor". Deficiency conditions may include hyperglycemia, transient hyper/hypoglycemia, fatigue, accelerated atherosclerogenesis, elevated LDL cholesterol, increased need for insulin and diabetes-like symptoms, and impaired stress responses. Marginal or insufficient Cr is common in the U.S., where average tissue levels are low compared to those found in many other countries. Low hair Cr appears to be associated with increased risk of cardiovascular disease and an atherogenic lipoprotein profile (low HDL, high LDL). Common causes of deficiency are ingestion of highly processed foods, inadequate soil levels of Cr, gastrointestinal dysfunction, and insufficient vitamin B-6. Cr status is also compromised in patients with iron overload/high transferrin saturation because transferrin is a major transport protein for Cr.

Confirmatory tests for Cr adequacy include glucose tolerance and packed red blood cell elements analysis.

Vanadium Low

Vanadium (V) is typically found at low levels in hair and the clinical significance of the measured result of lower than average hair V is not known. V is measured in hair for research purposes because it has been postulated to be an essential microtrace element. Indirect data to support this postulate have been derived from experimental models. Suggested functions for V include: regulation of sodium-potassium-ATPase, intracellular glutathione metabolism, thyroid metabolism, and insulin mimetic effects at pharmacological doses.

Average dietary V intake varies considerably between 20 mcg to 2 mg. Food sources of V include: liver, fish, radishes, grains, nuts, and vegetable oils.

Strontium High

Hair usually reflects the body burden of Strontium (Sr), and Sr levels usually correlate with calcium levels in body tissue. However, hair levels of Sr can be raised by external contamination, usually from hair treatment products. Elevated Sr in hair treated with permanent solutions, dyes, or bleaches is likely to be an artifact of hair treatment and probably does not reflect the level of Sr in other tissues.

Diseases of excess Sr have not been reported, except for Sr rickets. In general, Sr excess is not of clinical concern in the U.S. It's bad reputation comes from it's radioactive isotopes which were widespread in the western U.S. as a result of nuclear testing in the 1950's. Stable Sr (not radioactive Sr) is measured and reported by DDI.

Other tests indicative of Sr status or excess are measurements of Sr in whole blood, Sr/calcium ratio in blood, and Sr in urine.

Total Toxic Element Indication

The potentially toxic elements vary considerably with respect to their relative toxicities. The accumulation of more than one of the most toxic elements may have synergistic adverse effects, even if the level of each individual element is not strikingly high. Therefore, we present a total toxic element "score" which is estimated using a weighted average based upon relative toxicity. For example, the combined presence of lead and mercury will give a higher total score than that of the combination of silver and beryllium.

Lab number: **H160415-2375-1**
Patient: **Anna M. Salanti**

Hair

Page: 6
Client: **26339**
